

SURVEY OF FRESH WATER BODIES OF AJMER

**Dr Rashmi Sharma , Ashok Sharma , Amogh Bhardwaj and Devesh
Bhardwaj**

Associate Professor

SPCGCA

MDSU

AJMER

sharmarashmigca@gmail.com

Abstract

Ajmer is situated in the center of Rajasthan , also known as heart of Rajasthan . Rajasthan is the western arid state of INDIA. Ajmer is situated $25^{\circ} 38' - 26^{\circ} 56'$ north latitude and $73^{\circ} 54' - 75^{\circ} 22'$ East longitude. Area of Ajmer is 8481 sq km. Population of Ajmer is 5.43 lakhs. Ajmer has semiarid Eastern plane and Arid western plane. Semiarid and arid zones are divided by Aravallis , which are oldest mountains of the world.

Ajmer comes in Agroclimate IIIA zone. Ajmer has 9 tehsils and 8 Panchayat Samitis.

Total 1130 villages are there in Ajmer. Ajmer has 4 fresh water lakes , Anasagar , Foy sagar and

Pushkar and one more small Budha Pushkar. Ajmer gets its drinking water from Bisalpur dam

which is situated in Tonk district.

Introduction

The fresh water bodies of Ajmer are Anasagar , Foy sagar , Pushkar and Budha Pushkar :

Anasagar is feigned lake situated in center of Ajmer . It was made by

King Arnoraja (He was Grand father of King Prithviraj Chouhan). The lake spreads 13 km.

Baradari (Pavilion) was build by Shahjaha (1637) and the Garden Daulat Bagh by King

Jehangir . Circuit house was British Residency. The catchment area 1.9 sq mi (5 km), the depth of lake is 4.5 m the storage capacity of lake is 4750000 m³. (6210000cuyd.)

Foy sagar : Foy sagar is also unnatural lake it was named by Mr. Foy who created it during deprivation relief project (1892). Its capacity is 15 million cubic feet water spread 14,000,000 sq feet. If rainfall is low all water evaporates and Foy sagar artificial lake dries

up. This has become several times. Foy sagar is also endangered lake.

Pushkar : Pushkar is hallowed lake and the ingress of Thar desert. having 52 ghats.

Pilgrims take venerated dip on Purnima(Full moon day), Amavasya (No moon day), Kartik month (November) and Ekadashi (11th moon day), Dwadashi (12th moon day) and other

propitious occasions. It is situated at altitude of 510m.(Image :1).

Budha Pushkar : Budha Pushkar is positioned in Aravalli breach. It is surrounded by sand and

agriculture land. Budha Pushkar is located 25°25' – 26°35' North latitude and 74°37' – 74°42' East longitude. It is prime lake where Lord Ram performed last sacrament of King Dashrath here.

Holy Adyah temple is located here. Budha Pushkar is endangered (threatened) lake. Total volume of lake is 150 cu.m. water spread is 45 hectare. Ground water is 10 feet deep. It is eutropical and endangered lake.

Lokeshwari H, & G.T. Chandrappa (2006) studied impact of heavy metal contamination of Bellandur lake on soil and cultivated vegetation. Bhuvaneshwaran & N.G. Santhalakshmi and S. Rajeshwari (1999) studied water quality of river Adyar in Chennai city.

Lokeshwari H, & G.T. Chandrappa (2006) studied heavy metal content in water hyacinth & sediments of Lalbagh lake Bangalore. Laxen, D. and R.C. Harrison (1997). Studies the highway as a source of water pollution with heavy metal – lead. Kishore, M.A. & J.F. Machiwa. 2003.

Studied distribution of heavy metal in sediments.

Gopal , (1994) Studied conservation of inland water bodies of India. Zedler ,(1996) studied restoration requirement ,reconstruction of antecedent physical condition ,chemical adjustment of soil and water biological manipulation . Water quality of freshwater lakes of Rajasthan were studied in the present paper.

Methods

Survey was done to identify fresh water lakes of Ajmer.The samples were analysed with Standard methods (APHA 1989), Trivedy and Goel 1986.Zooplankton (Edmondsom 1959, Tonapi 1980. Water was collected in clean Plastic bottles of 2 liters .

Results And Discussion

Physiochemical parameters of Anasagar , Foysagar ,Pushkar and Budha Pushkar are given in Table 1.

Table 1. Physiochemical parameters of Anasagar , Foysagar, Pushkar and Budha Pushkar Ajmer.

S.No.(Parameter)	Anasagar	Foysagar	Pushkar	Buda Pushkar
1.Depth	4.4m	74 ft (22.5m)	8-10m	8 m
2.Area(spread)	13km	1400000sq feet 15 m cu feet	10 km	150m cu feet
3. Temperature ° C	18.2-29 ° C	17-30.2 ° C	15-39.0	15.0-39.6 °
4. pH	7.4-8.9	7.1-8.7	7.8	7.2-8.6
5. Conductivity			.32	.8-1.83
6. TDS mg/l	1400-1600	900	180	838.40-1171.20
7.DO mg/l	5.4-10	2.63-5.4	5.2	1.60-7.9
8.Alkalinity mg/l	1400-1700	90-109	50	292.3-775.00
9.Hardness mg/l			300-600	96.90-338.90
10.Ca hardness	88.5			79-258.
11.Mg hardness	250-474			12.2-157.7
12.Ca mg/l				23.5-105.9
13.Mg mg/l				2.9-157.7
14.Cl mg/l	430-590	25.5-75.6	39	31.2-98.5
15.Total solids	1487-1707	900		1330-2971.
16. TSS mg/l	23-26	211-900	180	448-2449.
17.Sodium mg/l	30		36	12.5-68.3
18. Potasium	17-24.7		3.6	8-43.7
19. Free Co2		1.5-5.5		4.5-47.5
20. Nitrate mg/l	10-17.9	.1-.38		.51-1.5

21.Phosphate mg/l	.11-1.7		.35	2.1-2.9
22.SO4			48 mg/l	
23.Flouride mg/l			0.2 mg/l	
24.BOD mg/l	3-10			
25.COD	10-41			
26.conductivity	2140-3500			
27.ColiformMPN/100ml	1110-8080			

Sample water of lakes were analyzed . The average pH of Anasagar lake was 8.2.

Foysagar lake pH 7.5. Pushkar lake pH was 7.7 and Budha Pushkar pH was 7.8.

pH of Anasagar lake was maximum. Other parameters studied were D. O. which was highest for

Anasagar lake. Alkalinity (1500-1600) was also highest for Anasagar lake. T.D.S. was also highest

for Anasagar lake 1500 -1600. Value of heavy metals Calcium , Magnisium sodium and hardness

were also high values for Anasagar lake. These may be due to Anasagar being in Ajmer city and

on roads nearby Anasagar vehicular discharge , agricultural runoff, sewage waste , during rains these all pollutants cause pollution of aquatic water.



Image :1 Pushkar Lake



Image 2 : Anasagar lake



Image 3 :Foisagar lake



Image 4: Budha Pushkar Lake

Conclusion and Summary

Anasagar ,Foy sagar , Pushkar and Buda Pushkar are the 4 main lakes of Ajmer ,of which Anasagar is largest. Eutrophication , increasing human population, Human activities ,quarrying in Aravallis ,religious ceremonies , tourists , low rainfall , high temperature , evanescence of water , urbanization , waste sewage is added to lake, detergenys, chemical waste, fertilizers, aquaculture, horticulture , urban setellement, pesticide waste, heavy metal waste, heavy metal waste are bioaccumulated and not good for human health.Cutting of trees are some more reasons for vandalization of lakes. These can be restored by planting trees, conservation of lakes, prohibiting human activities around lakes.

Allowing Religious Ceremonies at certain places only, ban of polythene. Pushkar and Budha Pushkar are surrounded by sand dunes . High temperature , low rainfall, high evaporation , high wind speed 45-65 k/hr, highest during May June. More dams should be made to

conserve water and solve water problem of Rajasthan. Interlinking of rivers is also required.

Acknowledgements : We are thankful to Principal SPCGCA for giving permission to do the work.

References

1. Arti D. Sharma , A and Antal , N. 2013 Zooplankton Diversity and Physio chemical condition of Temple pond in Birpur (J. & K ., India). International Research Journal of Environment Science.: 25 -30.
2. Abbasi , S. A. 1998 water quality sampling and Analysis 1 st Ed. Discovery Publishing House , New Delhi .
3. Ahamad , U. Parveen , S. Mola , H.R.A. Kabir , H.A. and Ganai A.H. 2012 .Zooplankton population in relation to Physiochemical parameters of Lal Diggi Pond in Aligarh , India j. Environ . Biol. 33: 1015-1019.
4. Benarjea S M 1967 Water quality and soil condition of fish ponds in some states of India In relation to fish production Indian Journal of Fisheries 14 (1 & 2) : 115-144.
5. Battish , S.K. 1992. Fresh water Zooplankton of India . Oxford and I B K Publishing Co. New Delhi.
6. Bhuvaneswaran , N. G. Santhalakshmi and S. Rajeshwari . 1999 . Water quality of Adyar in Chennai city . Indian Journal of Environment Protection 19 (6) : 412-415.
7. Kische , M.A. and J.F. Machiwa . 2003. Distribution of heavy metals in sediments of Mwanza Gulf Lake Victoria , Tanzania . Environmental International 28: 619-625.
8. Laxen , D. and R.C. Harrison . 1997 . The highways as a source of waterpollution as appraisal with the heavy metal - lead . Water Resources 11 : 1-11.
9. Lokeshwari , H. and G.T. Chandrappa . 2006. Heavy metals content in water , water hyacinth and sediments of lalbagh tank , Bangalore . Jr of Env. Sc & Engineering 48 (3) 183-188.
10. Lokeshwari , H. and G.T. Chandrappa . 2006. Impact of heavy metal contamination of

Bellandur lake on soil and cultivated vegetation . Current Science 91 (5) : 622-627.

11. Bordoloi , R., Abujam , S.K.S. and Paswan G. 2012. Assessment of fish culture in some fresh water ponds of Dhar Town , M.P., India ISCA Journal of Bio.

12.Gopal, 1994 . Conservation of inland water bodies of India.

13. Trivedy ,R.K. and Goel, P.K. 1986 . Chemical and Biological methods for water pollution studies.Environment Publications Karad , India.

14. Welch , P.S. 1952 . Limnology . Mc Graw Hill book Co. Inc., New York.

15.Tonapi ,G.T. 1980. Fresh water animals of India.Oxford and IBH publishingCompany ,New Delhi.

16.Sharma, M.S. 1980. Studies on Plankton and Productivity of Udaipur ,Water in Comparison to the selected Waters of Rajasthan .PhD. Thesis , University of Udaipur , India.

17.Zedler ,1996 .Restoration requirement reconstruction of antecedent physical condition , chemical adjustment of soil and water. Biological manipulation .