

Analysis of the tadpole graph $T_{m,n}$ cardinality of hop dominance number

¹M.Suneetha Rani, ²Sk.Jhansi Begum, ³B.Srinivasa Rao

^{1,2,3}Assistant Professor, Dept. of BS &H, Newton's Institute of Engineering, Macherla, Andhra Pradesh, India.

Abstract

Make $T_{m,n}$ a tadpole graph. If there is an u in S_h such that $d(u, v) = 2$ for all v in $V - S_h$, then the set $S_h \subseteq V(T_{m,n})$ is a hop dominating set of $T_{m,n}$. The hop domination number of G is the minimal cardinality of a hop dominating set of G and is represented by the symbol $h(T_{m,n})$. In this essay, we spoke about the tadpole graph's hop dominance number.

Keywords: hop-domination, hop-domination number, Tadpole graph, neighbourhood.

1. Introduction

[6] The graph created by connecting a cycle graph and a route graph with a bridge is known as the Tadpole graph (Truszczynski 1984) or Kite graph (Kim and Park 2006). $T_{m,n}$ serves as a sign for it. In specifically, $T_{3,1}$ and $T_{4,1}$ of the Tadpole graph ($T_{m,1}$) are referred to as the Paw graph and Banner graph, respectively. the graph of the generalized tadpole




Fig. 1.1

Let us denote the vertices of a Tadpole graph as two distinct sets:

- (i) Refer the vertices of the cycle graph C_m as $\{v_1, v_2, \dots, v_m\}$ and
- (ii) The Vertices of the Path graph P_n as $\{u_1, u_2, \dots, u_n\}$

\therefore The vertices of $T_{m,n}$ are

$$V(T_{m,n}) = V(C_m) \cup V(P_n) \\ = \{v_1, v_2, \dots, v_m, u_1, u_2, \dots, u_n\}$$








Theorem 1.1 ([11] p.546): A dominating set D of a graph G is minimal iff for each vertex $v \in D$, one of the following conditions satisfied,

- (i) There exists a vertex $u \in V - D$ such that $(u) \cap D = \{v\}$
- (ii) v is an isolated vertex in D .

[3] A subset S_h of $(T_{m,n})$ is a hop dominating set of $T_{m,n}$ if for all v in $V - S_h$, there exists u in S_h such that $(u, v) = 2$. The minimum cardinality of a hop dominating set of G is called the hop domination number of G

and is denoted by $\gamma_h(T_{m,n})$. For any vertex $v \in (T_{m,n})$, the open neighbourhood of v is the set $N(v) = \{u \in (T_{m,n}) | uv \in E(T_{m,n})\}$ and the closed neighbourhood is $N[v] = N(v) \cup \{v\}$. For a set $S_h \subseteq (T_{m,n})$, the open neighbourhood of S_h is $N(S_h) = \cup_{v \in S_h} N(v)$ and the closed neighbourhood is $N[S_h] = N(S_h) \cup S_h$. A set $S_h \subseteq (T_{m,n})$ is hop dominating set if $N[S_h] = V(T_{m,n})$.

2. Diagrammatic discussion on Hop domination number of Tadpole Graph

S.No.	Pan Graph (P_n)	Graph	$\gamma_h(G)$
1	$n=3, P_3$		2
2	$n=4, P_4$		2
3	$n=5, P_5$		2
4	$n=6, P_6$		3
5	$n=7, P_7$		3
6	$n=8, P_8$		4
7	$n=9, P_9$		4

8	$n=10, P_{10}$		4
---	----------------	--	---

Table 2.1: Pan Graph(P_n)

S.No.	Tadpole Graph ($T_{m,n}$), $m = 3$	Graph	$\gamma_h(G)$
1	$n=1, T_{3,1}$ (paw graph)		2
2	$n=2, T_{3,2}$		2
3	$n=3, T_{3,3}$		2
4	$n=4, T_{3,4}$		2
5	$n=5, T_{3,5}$		3
6	$n=6, T_{3,6}$		4
7	$n=7, T_{3,7}$		4
8	$n=8, T_{3,8}$		4










9	$n=9, T_{3,9}$		4
10	$n=10, T_{3,10}$		4

Table 2.2: Tadpole Graph (T_m), $m = 3$

S.No.	Tadpole Graph ($T_{m,n}$), $m = 4$	Graph	$\gamma_h(G)$
1	$n=1, T_{4,1}$ (Banner graph)		2
2	$n=2, T_{4,2}$		2
3	$n=3, T_{4,3}$		2
4	$n=4, T_{4,4}$		3
5	$n=5, T_{4,5}$		4
6	$n=6, T_{4,6}$		4
7	$n=7, T_{4,7}$		4









8	$n=8, T_{4,8}$		4
9	$n=9, T_{4,9}$		4
10	$n=10, T_{4,10}$		5

Table 2.3: Tadpole Graph (T_m), $m = 4$

S. No	Tadpole Graph ($T_{m,n}$), $m = 5$	Graph	$\gamma_h(G)$
1	$n=1, T_{5,1}$		2
2	$n=2, T_{5,2}$		2
3	$n=3, T_{5,3}$		2
4	$n=4, T_{5,4}$		3
5	$n=5, T_{5,5}$		4






6	$n=6, T_{5,6}$		4
7	$n=7, T_{5,7}$		4
8	$n=8, T_{5,8}$		4
9	$n=9, T_{5,9}$		4
10	$n=10, T_{5,10}$		5

Table 2.4: Tadpole Graph (T_m), $m = 5$

3. Results on Hop domination number of Tadpole graph $T_{m,n}$

Theorem 3.1: For m -pan graph, the hop domination number is given by

$$\gamma_h = \begin{cases} 2p & \text{iff } m = 6p \\ 2p + 1 & \text{iff } m = 6p + 1 \\ 2p + 2 & \text{iff } m = 6p + r, 2 \leq r \leq 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(1.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S'_h = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case(i): If $m=6p$.

Let $S_h = \{c_{6k-5}, c_{6k-4} | k = 1, 2, \dots, p\}$. → (1)

If $v = c_{6k-5}$, then atleast one vertex of $\{p_1, c_{m-1}, c_{6k-3}, c_{6k-7} | k = 1, 2, \dots, p\}$ is not hop dominating with any vertex in S'_h . If $v = c_{6k-4}$, then atleast one vertex of $\{c_m, c_{6k-6}, c_{6k-2} | k = 1, 2, \dots, p\}$ is not hop dominated by any vertex in S'_h . Therefore, S'_h is not a hop dominating set. Hence S_h is the minimum. Since for each $k, 1 \leq k \leq p$, there exists c_{6k-5}, c_{6k-4} in $|S_h| = 2p$. $\gamma_h(T_{m,1}) = 2p$ if $m = 6p$.

Conversely, If $\gamma_h(T_{m,1}) = 2p = |S_h|$, where S_h is given by an equation (1). Hence $V - S_h = \{c_{6k-2}, c_{6k-3}, c_{6k-6}, c_{6k-7}, p_1 \mid k = 1, 2, \dots, p\}$. $|V - S_h| = 4p + 1$. We know that $V = (V - S_h) \cup S_h$, therefore $|V| = 4p + 1 + 2p = 6p + 1$. Hence $m = 6p$.

Thus $\gamma_h(T_{m,1}) = 2p$ iff $m = 6p$.

Case(ii): If $m=6p+1$.

Let $S_h = \{c_{6k-5}, c_{6k-4}, c_{m-2} \mid k = 1, 2, \dots, p\}$.

If $v = c_{6k-5}$ or c_{6k-4} , the minimality of S_h follows from the above case(i) or else if $v = c_{m-2}$, there is no vertex in S_h' hop dominating with c_{m-2} . Hence S_h' is not a hop dominating set. Thus S_h is minimum and for each k , $1 \leq k \leq p$, there exists c_{6k-5}, c_{6k-4} in S_h and there exists c_{m-2} in S_h independent of k . Therefore $|S_h| = 2k + 1$. $\gamma_h(T_{m,1}) = 2p + 1$ if $m = 6p + 1$.

Case(iii): If $m=6p+2$.

Let $S_h = \{c_{6k-5}, c_{6k-4}, c_{m-2}, c_{m-3} \mid k = 1, 2, \dots, p\}$.

If $v = c_{6k-5}$ or c_{6k-4} , the minimality of S_h follows from the above case(i) or else if $v = c_{m-2}$ or c_{m-3} there is no vertex in S_h' hop dominating with c_{m-2} or c_{m-3} respectively. Hence S_h' is not a hop dominating set. Thus S_h is minimal hop dominating set and for each k , $1 \leq k \leq p$, there exists c_{6k-5}, c_{6k-4} in $V - S_h$ and there exists c_{m-2}, c_{m-3} in S_h independent of k . Therefore $|S_h| = 2k + 2$.

Case(iv): If $m=6p+3$.

Let $S_h = \{c_{6k-5}, c_{6k-4}, c_{m-2}, c_{m-3} \mid k = 1, 2, \dots, p\}$.

If $v = c_{6k-5}$ or c_{6k-4} or c_{m-2} , the minimality of S_h follows from the above case(i) and case (iii) or if $v = c_{m-3}$ there is no vertex in S_h' hop dominating c_{m-3} in $V - S_h'$. Therefore $|S_h| = 2k + 2$.

Case (v): If $m=6p+4$ and $6p+5$.

Let $S_h = \{c_{6k-5}, c_{6k-4} \mid k = 1, 2, \dots, p\}$. The minimality of S_h follows from case(i) and $|S_h| = 2(2p + 1) = 2p + 2$. Hence $\gamma_h(T_{m,1}) = 2p + 2$ if $m = 6p + r, 2 \leq r \leq 5$.

Let us indicate the vertices of T_m , as two sets first to refer the vertices of cycle graph C_m as $\{c_1, c_2, \dots, c_m\}$ and the second to refer the vertices of path graph P_n as $\{p_1, p_2, \dots, p_n\}$. So the vertices of T_m , is denoted as $V(T_{m,n}) = \{\{c_1, c_2, \dots, c_m\} \cup \{p_1, p_2, \dots, p_n\}\}$. Let the dominating set of T_m , be S_h .

Theorem 3.2: When $m=6p$, the hop domination of a tadpole graph T_m , is given by

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k & \text{if } n = 6k + r, 0 \leq r \leq 2 \\ 2p + 2k + 1 & \text{if } n = 6k + 3 \\ 2p + 2k + 2 & \text{if } n = 6k + r, r = 4, 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$

such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case(i): If $n=6k$

Let $S_h = \{c_{6q-2}, c_{6q-3}, p_{6s-2}, p_{6s-2} | q = 1, 2, \dots, m \text{ and } s = 1, 2, \dots, p\}$

If $v = c_{6q-2}$, then there is no vertex in S_h' hop dominating c_{6q-4} and c_{6q} . If $v = c_{6q-3}$, then there is no vertex in S_h' hop dominating c_{6q-5} and c_{6q-1} . If $v = c_{6s-2}$, then there is no vertex in S_h' hop dominating c_{6s-4} and c_{6s} . If $v = c_{6s-3}$, then there is no vertex in S_h' hop dominating c_{6s-1} and c_{6s-5} .

Thus S_h' is not minimal hop dominating set. Hence S_h is the minimal hop dominating set.

Case(ii): If $n=6k+1$

Let $S_h = \{c_{6q-5}, c_{6q-4}, p_{6s-1}, p_{6s-2} | q = 1, 2, \dots, m \text{ and } s = 1, 2, \dots, p\}$

If $v = c_{6q-5}$, there is no vertex in S_h' hop dominating c_{6q-3}, c_{6q-7} . In particular, If $q = 1$, there is no vertex in S_h' hop dominating c_{m-1}, c_3 and p_1 . If $v = c_{6q-4}$, there is no vertex in S_h' hop dominating c_{6q-2} and c_{6q-6} . In particular, If $q = 1$, there is no vertex in S_h' hop dominating c_m and c_4 . If $v = p_{6s-1}$, there is no vertex in S_h' hop dominating p_{6s-3} and p_{6s+1} . If $v = p_{6s-2}$, there is no vertex in S_h' hop dominating p_{6s-4} and p_{6s} .

Thus S_h' is not minimal hop dominating set. Hence S_h is the minimal hop dominating set.

Case(iii): $n=6k+2$

Let $S_h = \{c_{6q}, c_{6q-1}, p_{6s}, p_{6s-1} | q = 1, 2, \dots, m \text{ and } s = 1, 2, \dots, p\}$

(i) If $v = c_{6q}$, there is no vertex in S_h' hop dominating c_{6q+2} and c_{6q-2} . In particular, If $v = c_m$, there is no vertex in S_h' hop dominating c_2, c_{m-2} and p_2 . (ii) If $v = c_{6q-1}$, there is no vertex in S_h' hop dominating c_{6q+1} and c_{6q-3} . In particular, If $v = c_{m-1}$, there is no vertex in S_h' hop dominating c_1, c_{m-3} and p_1 . (iii) If $v = p_{6s}$, there is no vertex in S_h' hop dominating p_{6s-2} and p_{6s+2} . If $v = p_{6s-1}$, there is no vertex in S_h' hop dominating p_{6s-3} and p_{6s+1} .

In the above cases (i), (ii) and (iii) for each $q, 1 \leq q \leq m$, there exists c_i and c_{i+1} in S_h and for each $s, 1 \leq s \leq p$, there exists p_i and p_{i+1} in S_h , hence $|S_h| = 2p + 2k$.

Thus $\gamma_h(T_{m,n}) = 2p + 2k$ if $m = 6p$ and $n = 6k + r, 0 \leq r \leq 2$.

Case(iv): If $n=6k+3$.

Let $S_h = \{c_{6q}, c_{6q+1}, p_{6s}, p_{6s+1} | q = 0, 1, \dots, m \text{ and } s = 0, 1, 2, \dots, p\}$

(i): If $v = c_1$, there is no vertex in S_h' hop dominating c_3 . (ii): If $v = c_{6q}$, proof follows from (i) of case(iii). (iii): If $v = c_{6q+1}$, there is no vertex in S_h' hop dominating c_{6q-1}, c_{6q+3} . (iv): If $v = p_{6s}$, the proof follows from (iii) of case(iii). (v): If $v = p_{6s+1}$, there is no vertex in S_h' hop dominating p_{6s-1}, p_{6s+3} .

Hence S_h' is not minimum. Thus S_h is the hop dominating set and $|S_h| = 2p + 2k + 1$.

Thus $\gamma_h(T_{m,n}) = 2p + 2k + 1$ if $m = 6p$ and $n = 6k + 3$.

Case(v): If $n=6k+4$.

Let $S_h = \{c_1, c_2, c_{6q+1}, c_{6q+2}, p_{6s+1}, p_{6s+2} | q = 0, 1, \dots, m \text{ and } s = 0, 1, 2, \dots, k\}$

(i): If $v = c_1$, the proof follows from (i) of case(iv). (ii): If $v = c_2$, there is no vertex in S_h' hop dominating c_4 . (iii): If $v = c_{6q+1}$, the proof follows from (iii) of case(iv). (iv): If $v = c_{6q+2}$, there is no vertex in S_h' hop dominating c_{6q} and c_{6q+1} . (v): If $v = p_{6s+1}$, the proof follows from (v) of case(iv). (vi): If $v = p_{6s+2}$, there is no vertex in S_h' hop dominating p_{6s} and p_{6s+4} .

Case(vi): If $n=6k+5$

Let $S_h = \{c_{6q+2}, c_{6q+3}, p_{6s+2}, p_{6s+3} | q = 0, 1, \dots, m - 1 \text{ and } s = 0, 1, 2, \dots, k\}$

(i): If $v = c_{6q+2}$, there is no vertex in S_h' hop dominating c_{6q+4} and c_{6q} . (ii): If $v = c_{6q+3}$, there is no vertex in S_h' hop dominating c_{6q+5} and c_{6q+1} . (iii): If $v = p_{6s+2}$, the proof follows from (iv) of case(v). (iv): If $v = p_{6s+3}$, there is no vertex in S_h' hop dominating p_{6s+5}, p_{6s+1} .

In case (v) and (vi), S_h' is not minimum. Thus S_h is the hop dominating set and $|S_h| = 2p + 2k + 2$ if $m = 6p$ and $n = 6p + r, r = 4$ and 5 .

Thus $\gamma_h(T_{m,n}) = 2p + 2k + 2$ if $m = 6p$ and $n = 6p + r, r = 4$ and 5 .

Theorem 3.3: When $m = 6p + 1$, the hop domination of a tadpole graph, T_m , is given by,

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 1 & \text{if } n = 6k + r, 0 \leq r \leq 2 \\ 2p + 2k + 2 & \text{if } n = 6k + r, 3 \leq r \leq 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case(i): $n=6k$., Let $S_h = \{c_1, c_{6q-2}, c_{6q-1}, p_{6s-3}, p_{6s-2} | q = 1, 2, \dots, p \text{ \& } s = 1, 2, \dots, k\}$

If $v = c_1$, there is no vertex in S_h' hop dominating c_1 . If $v = c_{6q-2}$ or c_{6q-1} or p_{6s-3} or p_{6s-2} , the proof follows from case(i) of theorem (3.3). Thus S_h' is not minimal.

Case(ii): $n=6k+1$. Let $S_h = \{c_2, c_{6q-1}, c_{6q}, p_{6s-1}, p_{6s-2} | q = 1, 2, \dots, p \text{ \& } s = 1, 2, \dots, k\}$

If $v = c_2$, there is no vertex in S_h' hop dominating c_2 . If $v = c_{6q}$ or c_{6q-1} , the proof follows from case(iii) of theorem (3.3). If $v = p_{6s-3}$ or p_{6s-2} , the proof follows from case(ii) of theorem (3.3).

Therefore, S_h' is not minimal.

Case(iii): $n= 6k+2$. Let $S_h = \{c_1, c_{6q}, c_{6q+1}, p_{6s}, p_{6s-1} | q = 0, 1, 2, \dots, p \text{ \& } s = 1, 2, \dots, k\}$.

If $v = c_1$ or c_{6q} or c_{6q+1} or p_{6s} , the proof follows from case(iv) of theorem (3.3). If $v = p_{6s-1}$, the proof follows from subcase (iv) of case(iii) of theorem (3.3).

Therefore, S_h' is not minimal. Hence S_h is hop dominating set and $|S_h| = 2p + 2k + 1$ if $n = 6k + r, 0 \leq r \leq 2$.

Case(iv): $n=6k+3$. Let $S_h = \{c_3, c_4, c_{6q-4}, c_{6q-5}, p_{6s}, p_{6s+1} | q = 2, \dots, p + 1 \text{ \& } s = 0, 1, 2, \dots, k\}$.

If $v = c_3$ or c_4 , the proof follows from case(v) of theorem (3.3). If $v = c_{6q-4}$ or c_{6q-5} , the proof follows from case(iii) of theorem (3.3). If $v = p_{6s}$ or p_{6s+1} , the proof follows from case(iv) of theorem (3.3).

Case(v): $n=6k+4$. Let $S_h = \{c_{6q-3}, c_{6q-4}, p_{6s+1}, p_{6s+2} | q = 1, 2, \dots, p \text{ \& } s = 0, 1, 2, \dots, k\}$.

If $v = c_{6q-3}$, the minimality of D follows from case(i) of theorem (3.3) or else if $v = c_{6q-4}$, it follows from case (ii) of theorem(3.3). If $v = p_{6s+1}$ or p_{6s+2} , the minimality of S_h follows from case(iv) of theorem(3).

Case(vi): $n=6k+5$. Let $S_h = \{c_{6q-2}, c_{6q-3}, p_{6s+2}, p_{6s+3} | q = 1, 2, \dots, p \text{ \& } s = 0, 1, 2, \dots, k\}$.

If $v = c_{6q-2}$ or c_{6q-3} , the minimality of S_h follows from case(i) of theorem(3.3). If $v = p_{6s+2}$ or p_{6s+3} , the minimality of S_h follows from case(vi) of theorem (3.3).

Hence S_h is the hop dominating set and $|S_h| = 2p + 2k + 2$ if $n = 6k + r, 3 \leq r \leq 5$. Thus when $m=6p+1$, $\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 1 & \text{if } n = 6k + r, 0 \leq r \leq 2 \\ 2p + 2k + 2 & \text{if } n = 6k + r, 3 \leq r \leq 5. \end{cases}$

Theorem 3.4: When $m = 6p + 2$, the hop domination of T_m , is given

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 4 \\ 2p + 2k + 3 & \text{if } n = 6k + 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): If $n = 6k$, $S_h = \{c_1, c_2, c_{6q-1}, c_{6q}, p_{6s-3}, p_{6s-2} / q = 1, 2 \dots \dots, p \text{ \& } s = 1, 2, \dots \dots, k\}$

Case (ii): If $n = 6k + 1$, $S_h = \{c_2, c_3, c_{6q}, c_{6q+1}, p_{6s-2}, p_{6s-1} / q = 1, 2 \dots \dots, p \text{ \& } s = 1, 2, \dots \dots, k\}$

Case (iii): If $n = 6k + 2$, $S_h = \{c_3, c_4, c_{6q+1}, c_{6q+2}, p_{6s-1}, p_{6s} / q = 1, 2 \dots \dots, p \text{ \& } s = 1, 2, \dots \dots, k\}$

Case (iv): If $n = 6k + 3$, $S_h = \{c_{6q+2}, c_{6q+3}, p_{6s}, p_{6s+1} / q = 0, 1, \dots \dots, p \text{ \& } s = 0, 1, \dots \dots, k\}$

Case (v): If $n = 6k + 4$, $S_h = \{c_{6q-3}, c_{6q-2}, p_{6s+1}, p_{6s+2} / q = 1, 2 \dots \dots, p \text{ \& } s = 0, 1, 2, \dots \dots, k\}$

When $m = 6p + 2$, the minimality of S_h follows as the previous theorem and $|S_h| = 2p + 2k + 2$ if $n = 6k + r, 0 \leq r \leq 4$.

Case (vi): $n = 6k + 5, S_h = \{c_1, c_{6q-1}, c_{6q-2}, p_{6s+2}, p_{6s+3} / q = 1, 2 \dots \dots, p \ \& \ s = 0, 1, 2, \dots \dots, k\}$

When $m = 6p + 2$ and $n = 6k + 5$, the S_h is the hop dominating set as of from the previous theorems and $|S_h| = 2p + 2k + 3$.

Thus when $m = 6p + 2, \gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 4 \\ 2p + 2k + 3 & \text{if } n = 6k + 5. \end{cases}$

Theorem 3.5: When $m = 6p + 3$, the hop domination of T_m , is given

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 4 \\ 2p + 2k + 4 & \text{if } n = 6k + 5 \end{cases}$$

Proof:

Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): If $n = 6p, S_h = \{c_2, c_3, c_{6q}, c_{6q+1}, p_{6s-3}, p_{6s-2} / q = 1, 2 \dots \dots, p \ \& \ s = 1, 2, \dots \dots, k\}$

Case (ii): If $n = 6p + 1, S_h = \{c_{6q+1}, c_{6q+2}, p_{6s-2}, p_{6s-1} / q = 0, 1, 2 \dots \dots, p \ \& \ s = 1, 2, \dots \dots, k\}$

Case (iii): If $n = 6p + 2, S_h = \{c_{6q+2}, c_{6q+3}, p_{6s-1}, p_{6s} / q = 0, 1, 2 \dots \dots, p \ \& \ s = 1, 2, \dots \dots, k\}$

Case (iv): If $n = 6p + 3, S_h = \{c_{6q-3}, c_{6q-2}, p_{6s}, p_{6s+1} / q = 1, 2 \dots \dots, p \ \& \ s = 0, 1, \dots \dots, k\}$

Case (v): If $n = 6p + 4, S_h = \{c_{6q-2}, c_{6q-1}, p_{6s+1}, p_{6s+2} / q = 1, 2 \dots \dots, p \ \& \ s = 0, 1, 2, \dots \dots, k\}$

Case (vi): If $n = 6p + 5, S_h = \{c_1, c_2, c_{6q-1}, c_{6q}, p_{6s+2}, p_{6s+3} / q = 1, 2 \dots \dots, p \ \& \ s = 0, 1, 2, \dots \dots, k\}$

When $m = 6p + 3$, the minimality of S_h follows as theorem (2) & (3) and $|S_h| = \gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 4 \\ 2p + 2k + 4 & \text{if } n = 6k + 5. \end{cases}$

Theorem 3.6: When $m = 6p + 4$, the hop domination of a tadpole graph T_m , is given by

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 3 \\ 2p + 2k + 3 & \text{if } n = 6k + 4 \\ 2p + 2k + 4 & \text{if } n = 6k + 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): If $n = 6p, S_h = \{c_{6q+1}, c_{6q+2}, p_{6s-3}, p_{6s-2} / q = 0, 1, 2 \dots \dots, p \ \& \ s = 1, 2, \dots \dots, k\}$

Case (ii): If $n = 6p + 1, S_h = \{c_{6q+2}, c_{6q+3}, p_{6s-2}, p_{6s-1} / q = 0,1,2 \dots, p \ \& \ s = 1,2, \dots, k\}$

Case (iii): If $n = 6p + 2, S_h = \{c_{6q+3}, c_{6q+4}, p_{6s-1}, p_{6s} / q = 0,1,2 \dots, p \ \& \ s = 1,2, \dots, k\}$

Case (iv): If $n = 6p + 3, S_h = \{c_{6q+4}, c_{6q+5}, p_{6s}, p_{6s+1} / q = 0,1,2 \dots, p \ \& \ s = 1,2, \dots, k\}$

Case (v): If $n = 6p + 4, S_h = \{c_2, c_{6q-1}, c_{6q}, p_{6s+1}, p_{6s+2} / q = 1,2 \dots, p \ \& \ s = 0,1,2, \dots, k\}$

Case (vi): If $n = 6p + 5, S_h = \{c_1, c_2, c_{6q}, c_{6q+1}, p_{6s+1}, p_{6s+2} / q = 1,2 \dots, p \ \& \ s = 0,1, \dots, k\}$

When $m=6p+4$, the minimality of S_h follows as in theorem (3.2) & (3.3) and

$$|S_h| = \gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 3 \\ 2p + 2k + 3 & \text{if } n = 6k + 4 \\ 2p + 2k + 4 & \text{if } n = 6k + 5. \end{cases}$$

Theorem 3.7: When $m = 6p + 5$, the hop domination of T_m , is given by

$$\gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 3 \\ 2p + 2k + 4 & \text{if } n = 6k + r, r = 4 \text{ and } 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): If $n = 6p, S_h = \{c_{6q+2}, c_{6q+3}, p_{6s-3}, p_{6s-2} / q = 0,1,2 \dots, p \ \& \ s = 1,2, \dots, k\}$

Case (ii): If $n = 6p + 1, S_h = \{c_{6q+3}, c_{6q+4}, p_{6s-2}, p_{6s-1} / q = 0,1,2 \dots, p \ \& \ s = 1,2 \dots, k\}$

Case (iii): If $n = 6p + 2, S_h = \{c_{6q+4}, c_{6q+5}, p_{6s-1}, p_{6s} / q = 0,1,2 \dots, p \ \& \ s = 1,2 \dots, k\}$

Case (iv): If $n = 6p + 3, S_h = \{c_{6q}, c_{6q-1}, p_{6s}, p_{6s+1} / q = 1,2 \dots, p \ \& \ s = 0,1,2 \dots, k\}$

Case (v): If $n = 6p + 4, S_h = \{c_2, c_3, c_{6q}, c_{6q+1}, p_{6s+1}, p_{6s+2} / q = 1,2 \dots, p \ \& \ s = 0,1,2 \dots, k\}$

Case (vi): If $n = 6p + 5, S_h = \{c_{6q+3}, c_{6q+4}, p_{6s+2}, p_{6s+3} / q = 0,1,2 \dots, p \ \& \ s = 0,1,2 \dots, k\}$

When $m = 6p + 5$, the minimality of S_h follows as in theorem (2) and (3), Hence

$$|S_h| = \gamma_h(T_{m,n}) = \begin{cases} 2p + 2k + 2 & \text{if } n = 6k + r, 0 \leq r \leq 3 \\ 2p + 2k + 4 & \text{if } n = 6k + r, r = 4 \text{ and } 5. \end{cases}$$

Theorem 3.8: For $n = 2$ the hop domination of a Tadpole graph T_m , is given by

$$\gamma_h(T_{m,2}) = \begin{cases} 2p & \text{if } m = 6p \\ 2p + 1 & \text{if } m = 6p + 1 \\ 2p + 2 & \text{if } m = 6p + r, 2 \leq r \leq 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): $m = 6p$, Let $S_h = \{c_{6q}, c_{6q-1} / q = 1, 2, \dots, k\}$

- (a) If $v = c_{6q}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_2, p_2, c_{6q+2}, c_{6q-2}\}$.
 (b) If $v = c_{6q-1}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_1, p_1, c_{6q+1}, c_{6q-3}\}$

Thus S_h' is not minimum. Hence S_h is the hop dominating set and $|S_h| = 2p$.

Case (ii): $m = 6p + 1$, Let $S_h = \{c_3, c_{6q}, c_{6q+1} / q = 1, 2, \dots, k\}$

- (a) If $v = c_3$, \exists no vertex in S_h' hop dominating c_3 . (b) If $v = c_{6q}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_1, p_1, c_{6q+2}, c_{6q-2}\}$. (c) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_2, p_2, c_{6q+3}, c_{6q-1}\}$.

Thus S_h is the hop dominating set and $|S_h| = 2p + 1$.

Case (iii): $m = 6p + 2$, Let $S_h = \{c_{6q+1}, c_{6q+2} / q = 0, 1, \dots, k\}$.

- (a) If $v = c_{6q+2}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q+1}, c_{6q-1}$ or $p_1\}$.
 (b) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q}, c_{6q+4}$ or $p_2\}$

Thus S_h is the hop dominating set and $|S_h| = 2p + 2$

Case (iv): $m = 6p + 3$, Let $S_h = \{c_{6q+2}, c_{6q+3} / q = 0, 1, \dots, k\}$

- (a) If $v = c_{6q+2}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q}, c_{6q+4}, p_1\}$.
 (b) If $v = c_{6q+3}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q-1}, c_{6q+5}, p_2\}$.

Thus S_h is the hop dominating set and $|S_h| = 2p + 2$

Case (v): $m = 6p + 4$, Let $S_h = \{c_{6q+3}, c_{6q+4} / q = 0, 1, \dots, k\}$

- (a) If $v = c_{6q+3}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q-1}, c_{6q+5}, p_1\}$.
 (b) If $v = c_{6q+4}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q-2}, c_{6q+6}, p_2\}$.

Thus S_h is the hop dominating set and $|S_h| = 2p + 2$.

Case (vi): $m = 6p + 5$, Let $S_h = \{c_{6q+4}, \frac{c_{6q+5}}{q} / q = 0, 1, \dots, k\}$

- (a) If $v = c_{6q+4}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q-2}, c_{6q+6}, p_1\}$.
 (b) If $v = c_{6q+5}$, \exists no vertex in S_h' hop dominating with at least one of the vertex of $\{c_{6q-3}, c_{6q+7}, p_2\}$.

Thus S_h is the hop dominating set and $|S_h| = 2p + 2$

$$\text{Thus, } h(T_{m,2}) = \begin{cases} 2p & \text{if } m = 6p \\ 2p + 1 & \text{if } m = 6p + 1 \\ 2p + 2 & \text{if } m = 6p + r, 2 \leq r \leq 5 \end{cases}$$

Theorem 3.9: For $n = 3$, the hop domination of a Tadpole graph T_m , is given by

$$\gamma_h(T_{m,3}) = \begin{cases} 2p + 1 & \text{if } m = 6p \\ 2p + 2 & \text{if } m = 6p + r, 1 \leq r \leq 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): $m = 6p$, Let $S_h = \{c_{6q}, c_{6q+1}, p_1 / q = 0, 1, 2, \dots, k\}$

(a) If $v = c_{6q}$, \exists no vertex in S_h' hop dominating with atleast one of the vertices such as of $\{c_2, c_{6q+2}, c_{6q-2}, p_2\}$. (b) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating with atleast one of the vertices such as of $\{c_{6q-3}$ and $c_{6q-1}\}$. (c) If $v = p_3$, \exists no vertex in S_h' hop dominating with atleast one of the vertices such as of $\{c_1, c_{m-1}$ and $p_3\}$

Thus S_h is the minimal hop dominating set and $|S_h| = 2p + 1$

Case (ii): $m = 6p + 1$, Let $S_h = \{c_{6q}, c_{6q+1}, p_1 / q = 0, 2, \dots, k\}$

(a) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating with atleast one of the vertex such as of $\{c_{6q-1}, c_{6q+1}, c_2, p_2\}$. (b) If $v = c_{6q+2}$, \exists no vertex in S_h' hop dominating with atleast one of the vertices such as of $\{c_{6q}$ and $c_{6q+4}\}$. (c) If $v = p_1$, \exists no vertex in S_h' hop dominating with atleast one of the vertices such as of $\{c_1, c_{m-1}$ and $p_3\}$

Thus S_h is the minimal hop dominating set and $|D| = 2p + 2$

Case (iii): $m = 6p + 2$, Let $S_h = \{c_{6q+2}, c_{6q+3}, p_1 / q = 0, 1, 2, \dots, k\}$

Case (iv): $m = 6p + 3$, Let $S_h = \{c_{6q+3}, c_{6q+4}, p_1 / q = 0, 1, 2, \dots, k\}$

Case (v): $m = 6p + 4$, Let $S_h = \{c_{6q+4}, c_{6q+5}, p_1 / q = 0, 1, 2, \dots, k\}$

Case (vi): $m = 6p + 5$, Let $S_h = \{c_{6q}, c_{6q+1}, p_1 / q = 0, 1, 2, \dots, k\}$

The Proof of case (iii) to (vi) follows same as previous cases. Thus S_h is the minimal hop-dominating set and $|S_h| = 2p + 2$

$$\text{Thus for } n = 3, \gamma_h(T_{m,3}) = \begin{cases} 2p + 1 & \text{if } m = 6p \\ 2p + 2 & \text{if } m = 6p + r, 1 \leq r \leq 5 \end{cases}$$

Theorem 3.10: For $n = 4$, the hop domination of a Tadpole graph T_m , is given by

$$\gamma_h(T_{m,4}) = \begin{cases} 2p + 2 & \text{if } m = 6p + r, 0 \leq r \leq 3 \\ 2p + 3 & \text{if } m = 6p + 4 \\ 2p + 4 & \text{if } m = 6p + 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such

that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): $m = 6p$, Let $S_h = \{c_{6q+1}, c_{6q+2}, p_1, p_2 / q = 0, 1, \dots, k - 1\}$

(a) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating $\{c_{6q-1}$ and $c_{6q+3}\}$. (b) If $v = c_{6q+2}$, \exists no vertex in S_h' hop dominating $\{c_{6q}$ and $c_{6q+4}\}$. (c) If $v = p_1$, \exists no vertex in S_h' hop dominating with $\{p_3\}$. (d) If $v = p_2$, \exists no vertex in S_h' hop dominating with $\{p_4\}$

Thus S_h is the minimal hop dominating set and $|S_h| = 2k + 2$

Case (ii): $m = 6p + 1$, Let $S_h = \{c_{6q+4}, c_{6q+5}, p_1, p_2 / q = 0, 1, \dots, k - 1\}$

The Proof is similar to case (i) and $|D| = 2p + 2$

Case (iii): $m = 6p + 2$, Let $S_h = \{c_{6q+3}, c_{6q+4}, p_1, p_2 / q = 0, 1, 2, \dots, k - 1\}$

The Proof is similar to case (i) and $|S_h| = 2p + 2$

Case (iv): $m = 6p + 3$, Let $S_h = \{c_{6q+4}, c_{6q+5}, p_1, p_2 / q = 0, 1, 2, \dots, k - 1\}$

The Proof is similar to case (i) and $|S_h| = 2p + 2$

Case (v): $m = 6p + 4$, Let $S_h = \{c_2, c_{6q-1}, c_{6q}, p_1, p_2 / q = 1, 2, \dots, k\}$

(a) If $v = c_2$, \exists no vertex in S_h' hop dominating $\{c_2\}$. (b) If $v = c_{6q-1}$, \exists no vertex in S_h' hop dominating $\{c_{6q-3}$ and $c_{6q+1}\}$. (c) If $v = c_{6q}$, \exists no vertex in S_h' hop dominating $\{c_{6q-2}$ and $c_{6q+2}\}$. (d) If $v = p_1$, \exists no vertex in S_h' hop dominating with $\{p_3\}$. (e) If $v = p_2$, \exists no vertex in S_h' hop dominating with $\{p_4\}$

Thus S_h is the minimal hop dominating set and $|S_h| = 2p + 3$

Case (vi): $m = 6p + 5$, Let $S_h = \{c_2, c_3, c_{6q}, c_{6q+1}, p_1, p_2 / q = 1, 2, \dots, k\}$

(a) If $v = c_2$, \exists no vertex in S_h' hop dominating $\{c_2\}$. (b) If $v = c_3$, \exists no vertex in S_h' hop dominating $\{c_3\}$. (c) If $v = c_{6q}$, \exists no vertex in S_h' hop dominating $\{c_{6q-2}$ and $c_{6q+2}\}$. (d) If $v = c_{6q+1}$, \exists no vertex in S_h' hop dominating $\{c_{6q-1}$ and $c_{6q+3}\}$. (e) If $v = p_1$, \exists no vertex in S_h' hop dominating with $\{p_3\}$. (f) If $v = p_2$, \exists no vertex in S_h' hop dominating with $\{p_4\}$

Thus S_h is the minimal hop dominating set and $|S_h| = 2p + 4$

$$\text{Hence when } n = 4, \gamma_h(T_{m,4}) = \begin{cases} 2p + 2 & \text{if } m = 6p + r, 0 \leq r \leq 3 \\ 2p + 3 & \text{if } m = 6p + 4 \\ 2p + 4 & \text{if } m = 6p + 5 \end{cases}$$

Theorem 3.11: For $n = 5$, the hop domination of a Tadpole graph T_m , is given by

$$\gamma_h(T_{m,4}) = \begin{cases} 2p + 2 & \text{if } m = 6p + r, r = 0, 1 \\ 2p + 3 & \text{if } m = 6p + 2 \\ 2p + 5 & \text{if } m = 6p + r, 3 \leq r \leq 5 \end{cases}$$

Proof: Let S_h be the hop dominating set of $T_{m,1}$. The minimality of S_h follows from theorem(3.1) using the contrary of this theorem. If S_h is not a minimal hop dominating set then there exists $v \in S_h$ such that $S_h' = S_h - \{v\}$ is a hop dominating set of $T_{m,1}$. Therefore for all $u \in N'[v]$ there exists $v' \in u \in N'[v] - \{v\}, v' \in N'[v]$.

Case (i): If $m = 6p$, Let $S_h = \{c_{6q+2}, c_{6q+3}, p_2, p_3 / q = 0, 1, \dots k - 1\}$

Case (ii): If $m = 6p + 1$, Let $S_h = \{c_{6q+3}, c_{6q+4}, p_2, p_3 / q = 0, 1, 2, \dots k - 1\}$

Case (iii): If $m = 6p + 2$, Let $S_h = \{c_{6q+4}, c_{6q+5}, p_2, p_3 / q = 0, 1, 2, \dots k - 1\}$

Case (iv): If $m = 6p + 3$, Let $S_h = \{c_3, c_{6q-1}, c_{6q}, p_2, p_3 / q = 1, 2, \dots k\}$

Case (v): If $m = 6p + 4$, Let $S_h = \{c_2, c_3, c_{6q}, c_{6q+1}, p_2, p_3 / q = 1, 2, \dots k\}$




Case (vi): If $m = 6p + 5$, Let $S_h = \{c_{6q+1}, c_{6q+2}, p_2, p_3 / q = 0, 1, \dots k\}$. The proof follows as the previous theorem.


Theorem 3.12: $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m})$ iff $m = n$.

Proof: Let $m = n$, then T_m and $T_{n,m}$ are same graphs. Hence $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m})$. On the other hand, assume $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m})$, suppose if $m \neq n$, then by theorems(--) $\gamma_h(T_{m,n}) \neq \gamma_h(T_{n,m})$. Thus $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m})$ iff $m = n$.

Corollary: $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m}) = m - 1$ (or) $n - 1$ iff $m = n$, where $m = n = 3, 4, 5$. and $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m}) \leq m - 1$ (or) $n - 1$ iff $m = n$, where $m = n = 6$.

Proof:

S.No.	Tadpole Graph $(T_{m,n}), m = n$	Graph	$\gamma_h(G)$
1	$m=n=3, T_{3,3}$		2 or (n-1)
2	$m=n=4, T_{4,4}$		3 or (n-1)
3	$m=n=5, T_{5,5}$		4 or (n-1)

4	$m=n=6, T_{6,6}$		4 or $\leq(n-1)$
---	------------------	---	------------------

From the table we get $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m}) = m - 1$ (or) $n - 1$ iff $m = n$, where

$m = n = 3,4,5$. And $\gamma_h(T_{m,n}) = \gamma_h(T_{n,m}) \leq m - 1$ (or) $n - 1$ iff $m = n$, where $m = n = 6$.

Conclusion

In this paper, we have found the hop domination number of Tadpole graph and derived some theorems on it.

References

- [1]. Harary, F., Graph Theory, ed. 1, Addison-Welsley, 1969.
- [2]. S.Alikhani and Y.H.Peng, *Dominating sets and Domination polynomials of Paths*, International Journal of Mathematics and Mathematical Science, (2009).
- [3]. C.Natarajan, S.K.Ayyaswamy and G.Sathiamoorthy, *A note on hop domination number of some special families of graphs*, International Journal of Pure and Applied Mathematics Vol 119, No.12, 2018, 14165-14171.
- [4]. M.Al-Harere and P.A. Khuda Bakhsh, *Changes of tadpole domination number upon changing of graphs*, Sci.Int.(Lahore), 31(2), 197-199, 2019. ISSN 1013-5316; CODEN: SINTE 8.
- [5]. A.Vijayan and T.Nagarajan, *Vertex-Edge Domination Polynomials of Lollipop Graphs $L_{n,1}$* , International Journal of Scientific and Innovative Mathematical Research (IJSIMR), Vol.-3, Issue 4, April 2015, PP 39-44, ISSN 2347-307X(Print) & ISSN 2347-3142(Online).
- [6]. Ayhan A.Khalil and Omar A.Khalil, *Determination and Testing the Domination numbers of Tadpole Graph, Book Graph and Stacked Book Graph Using MATLAB*, College of Basic Education Researchers Journal Vol.10, No.1.
- [7]. Nigar Siddiqui, Mohit James, *Domijnation and Chromatic Number of Pan Graph and Lollipop Graph*, International Journal of Technical Innovation in Modern Engineering & Science (IJTIMES), Impact factor: 5.22 (SJIF-2017), e-ISSN: 2455-2585, Vol-4, Issue 6, June-2018.
- [8]. M.S.Franklin Tamil Selvi and A.Amutha, *A Performance on Harmonious Coloring of Barbell Graph*, International Journal of Engineering and Advanced Technology (IJEAT), ISSN: 2249-8958, Vol.-9, Issue-1, October 2019.
- [9]. S.Nagarajan, Aswini.B and Vijaya.A, *Upper Bound of Hop Domination number for Regular Graphs of Even degree*, ANVESAK, ISSN: 0378-4568, Vol. 51, No.1(X) January-June 2021.
- [10]. S.Nagarajan, Vijaya.A and Aswini.B, *Lower Bound of Hop Domination Number for Regular Graphs of Odd degree*, ANVESAK, ISSN: 0378-4568, Vol. 51, No.1(X) January-June 2021.
- [11]. Kulli.V.R. and Janakiram.B, *The Nonsplit Domination Number of A Graph*, Indian J.Pure Appl.Math. Vol.31, No.5, pp.545-550.